



UTAH SOCIETY OF
HEALTH-SYSTEM PHARMACISTS

Annual Meeting
August 18, 2018

Poll Everywhere Audience Response

ACPE requires active learning and most prefer real-time participation rather than a graded post-test

We are utilizing Poll Everywhere software for this process.

You may join to participate by 3 different ways:

Text Messaging: Text ushp to 22333

Web Browser: Go to PollEv.com/ushp

Poll Everywhere app: Download app and join ushp presentation

For each question, you can click on the correct answer in Web Browser or App or text correct answer to 22333

Updated IDSA Guidelines for the Treatment of *C. difficile* Infection

Brian Hathaway, PharmD
Infectious Disease Pharmacist
St. Mark's Hospital



Disclosure

- I have no conflicts of interest to disclose.
- I will discuss the use of Fecal Microbiota Transplantation for the treatment of *C. difficile* infection.
- I will also discuss the dosing of rifaximin (Xifaxan[®]) in pediatric patients.
- No other off-label drug uses



Learning Objectives

At the conclusion of this activity, **pharmacists** should be able to successfully:

1. Evaluate and contrast new recommendations for drug treatment of *C. difficile* infection, particularly related to metronidazole and fidaxomicin.
2. Summarize the evidence supporting those changes, with an emphasis on long-term outcomes.
3. Formulate and defend a treatment recommendation to a provider, using the best available evidence from the new guidelines.
4. Teach a patient about *C. difficile* risk factors, therapy options and possible results.



Learning Objectives

At the conclusion of this activity, ***pharmacy technicians*** should be able to successfully:

1. Describe the mechanism and impact of *C. difficile* infection.
2. Identify medications in current use for the treatment of *C. difficile*.
3. Explain why antibiotic use is the most important modifiable risk in developing this infection.



Why should I care about *C. difficile*?

- Estimated annual incidence in US: 453,000 in 2011; 29,000 deaths
- Since 2000, CDI-attributable mortality: 4.5%-5.7% (endemic), 6.9%-16.7% (epidemic)
- Per episode, for acute-care hospitals: \$3427-\$9960 (2012 dollars)
- Extrapolates to \$1.2-\$5.9 billion annual acute care costs



Preliminaries



Abbreviations

- Vancomycin (Vancocin[®]) = Vancomycin (Vancocin[®])
PO/PR
- CDI: *Clostridium difficile* infection
- FMT: Fecal Microbiota Transplantation (sometimes
Fecal Matter Transplant)

Structure of Guidelines

1. Epidemiology

2. Diagnosis

3. Infection Prevention

1. Antibiotic Stewardship

2. PPIs and Probiotics

4. Treatment

1. Adults

2. Pediatrics

5. Unanswered Questions




Background

- Since 2016, ~~*Clostridium difficile*~~
 - NOW *Clostridioides difficile*
- Why the change?
 - Improved sequencing
 - Phenotype/biochemistry
- *C. difficile* colitis first described in 1970's
 - Broad-spectrum agents
 - Disruption of normal bowel flora



Bacterial clones in a random stool sample Before, during and after Augmentin




Toxigenic *C. difficile*


- Toxin A
 - Inflammation →
 - Fluid secretion →
 - Mucosal injury
- Toxin B
 - Opens cell junctions
 - Essential to toxigenic activity
- NAP1/BI/027
 1. “Hypervirulent”
 2. Binary toxin
 3. Increase in toxin production
 4. Fluoroquinolone resistance



Pathophysiology



Imaging



Clinical Diagnosis

- Which patients to test?
- Adults
 1. Unexplained
 2. New-onset
 3. ≥ 3 unformed stools in 24 hrs
- Pediatrics
 - ≤ 12 mos old = DO NOT TEST
 - 1-2 yrs = only if other causes for diarrhea ruled out
 - > 2 yrs = prolonged/worsening diarrhea, contact with healthcare

AND

AND



Question 1

Choose the toxin that is NOT seen in *C. difficile* infections.

A. Binary toxin

B. Toxin B

C. Tetrodotoxin

D. Toxin A



Laboratory Methods




Detection Techniques

- GDH enzyme: glutamate \rightarrow α -ketoglutarate + ammonia
- EIA – Enzyme Immunoassay
- NAAT – Nucleic Acid Amplification Test

- Is *C. difficile* here?
- Is *C. difficile* producing toxin?
- Is *C. difficile* producing LOTS of toxin?




Detection Techniques



Detection Algorithms

- What tests should we use?
 - Single step: NAAT
 - Multistep: GDH + toxin, GDH + toxin, mediated by NAAT, NAAT + toxin
- Multi-step algorithm recommended



Repeat Testing

- Adult and pediatric patients
 - Within 7 days of original episode: No
 - Asymptomatic patients: No
- Confounders
 - Asymptomatic carriage
 - Bacterial shedding



Infection Prevention



Infection Prevention Specifics

- Topics
 - Isolation
 - Personal protective equipment
 - Device and room cleaning
- Hand hygiene
 - Soap and water over alcohol



Stewardship is Good Practice

[It] is sufficiently obvious that desirable effects outweigh undesirable effects that no direct evidence is available because no one would be foolish enough to conduct a study addressing the implicit clinical question. Typically, such recommendations are supported by a great deal of indirect evidence, but teasing out the nature of the indirect evidence would be challenging and a waste of time and energy.



Stewardship Ideas

■ Targets

- Fluoroquinolones
- Clindamycin
- Cephalosporins
- Broad-spectrum agents

■ Methods

- Formulary restriction
- Prospective audit and feedback
- Use and duration



PPIs and Probiotics

- Proton pump inhibitors / histamine H₂-receptor blockers
 - Epidemiologic association
 - Surrogate markers for disease severity?
 - No recommendation

- Probiotics
 - Inconsistent data
 - No recommendation



Question 2

Antimicrobial stewardship of which classes would be most helpful in controlling CDI?

Choose all that apply

- A. Aminoglycosides
- B. Fluoroquinolones
- C. Streptogramins
- D. Cephalosporins



Treatment



Definitions

- Non-severe
 - Leukocytosis with white blood cell count <15,000 cells/mL
 - Serum creatinine <1.5 mg/dL

- Severe
 - Leukocytosis with white blood cell count >15,000 cells/mL

OR

 - Serum creatinine ≥1.5 mg/dL

- Fulminant
 - Hypotension or shock, ileus, megacolon



Initial Episode - Adult

- Stop the causative agent!
 - If possible, choose an alternative.

- Non-severe or Severe

1. Oral vancomycin (Vancocin®) 125 mg PO QID x 10 days
2. Fidaxomicin (Dificid®) 200 mg PO BID x 10 days

- No preference

- Non-severe only

- Metronidazole (Flagyl®) 500 mg PO Q8H
- Access is the question



Initial Episode - Adult

- Why move away from metronidazole (Flagyl[®])?
 - No trials comparing metronidazole (Flagyl[®]) and fidaxomicin (Dificid[®])!
- Direct comparisons
 - Metronidazole (Flagyl[®]) and vancomycin (Vancocin[®])
 - Vancomycin (Vancocin[®]) and fidaxomicin (Dificid[®])



Initial Episode - Adult

Outcomes	# Participants	% Resolution	Relative Effect (95% CI)	p Value	Quality of Evidence (GRADE)
Direct comparison of fidaxomicin (Difcid®) and vancomycin (Vancocin®)					
Resolution of diarrhea at the end of treatment	1105	88 (FDX) 86 (VAN)	RR, 0.9 (0.98-1.1)	0.36	High
Resolution of diarrhea at the end of treatment without CDI recurrence ~1 month after treatment	1105	71 (FDX) 57 (VAN)	RR, 1.2 (1.1-1.4)	<0.0001	High
Direct comparison of metronidazole (Flagyl®) and vancomycin (Vancocin®)					
Resolution of diarrhea at the end of treatment without CDI recurrence ~1 month after treatment	843	63 (MTR) 73 (VAN)	RR, 0.87 (0.79-0.96)	0.003	High

FDX = fidaxomicin (Difcid®)
MTR = metronidazole (Flagyl®)
VAN = vancomycin (Vancocin®)

Summary of all available trials published.
Conducted by guideline preparers.



Fulminant/Recurrent - Adult

Oral vancomycin (Vancocin®) 500 mg PO QID x 10 days

CONSIDER

Ileus: vancomycin (Vancocin®) 500 mg PR QID x 10 days

PLUS

Metronidazole (Flagyl®) 500 mg IV Q8H x 10 days

Surgical management is definitive therapy.



1st Recurrence - Adult

Oral vancomycin (Vancocin[®]) tapered and pulsed
125 mg PO QID x 10 days, BID x 7 days, daily x 7 days, every 2-3 days for 2-8 weeks

OR

Fidaxomicin (Difacid[®]) 200 mg PO BID x 10 days instead of oral vancomycin (Vancocin[®]) x 10 days

OR

If metronidazole (Flagyl[®]) initially, then oral vancomycin (Vancocin[®]) x 10 days



≥ 2nd Recurrence - Adult

Vancomycin (Vancocin[®]) tapered and pulsed

OR

Vancomycin (Vancocin[®]) 125 mg PO QID x 10 days, followed by rifaximin (Xifaxan[®]) 400 mg PO TID x 20 days

OR

Fidaxomicin (Dificid[®]) 200 mg PO BID x 10 days

OR

Fecal Microbiota Transplantation (FMT)



>1 Recurrence – Adult and Pediatric




Fecal Microbiota Transplantation




Fecal Microbiota Transplantation

Increased diversity


Marked change in proportions



Fecal Microbiota Transplantation



Fecal Microbiota Transplantation



Question 3

A hospitalist asks you for a discharge recommendation for treatment of his patient's CDI. The patient is completing a 7-day course of levofloxacin for a UTI. She contracted the infection during this admission, and has never had *C. difficile* before. If needed, she can pay for PO vancomycin (Vancocin®). However, she would prefer to use a cheaper alternative if possible. What would you recommend?

- A. PO metronidazole (Flagyl®)
- B. PO vancomycin (Vancocin®)
- C. PO vancomycin (Vancocin®), only after failing PO metronidazole (Flagyl®)
- D. PO fidaxomicin (Difacid®)



Initial Episode - Pediatric

- Non-severe
 1. Oral vancomycin (Vancocin®) 10 mg/kg/dose PO QID x 7-10 days

OR

 2. Metronidazole (Flagyl®) 7.5 mg/kg/dose PO TID-QID x 10 days
 - No preference
- Severe/Fulminant
 1. Oral vancomycin (Vancocin®) PO/PR 10 mg/kg/dose PO QID x 7-10 days

AND / OR

 2. Metronidazole (Flagyl®) IV TID-QID x 10 days



1st Recurrence - Pediatric

Vancomycin (Vancocin[®]) PO tapered and pulsed (up to 12 weeks)

OR

Metronidazole (Flagyl[®]) PO x 10 days

No preference



≥ 2nd Recurrence - Pediatric

Vancomycin (Vancocin®) PO tapered and pulsed (up to 12 weeks)

OR

Vancomycin (Vancocin®) 125 mg PO QID x 10 days, followed by rifaximin (Xifaxan®) 400 mg PO TID x 20 days

OR

Fecal Microbiota Transplantation (FMT)



Question 4

A 9 year-old boy with recurrent CDI is admitted to your floor. While doing the medication reconciliation, you find that his parents are frustrated with this readmission. He has already taken courses of vancomycin (Vancocin[®]) and metronidazole (Flagyl[®]).

They ask you about alternatives to drug therapy, like “Poo in a Pill”. What options would you discuss?

- A. FMT is first-line therapy and more likely to clear CDI than other interventions.
- B. FMT could be reasonable, but only if antibiotics, probiotics and surgery have failed.
- C. FMT is not preferred, but is helpful in some patients, such as the immunosuppressed.
- D. FMT can be effective, used only after antibiotic therapy fails.



Conclusions

- Upgrades
 - Clarity of roles of fidaxomicin (Dificid®) and metronidazole (Flagyl®)
 - Recommendations for pediatric patients
 - Highlights role for antibiotic stewardship
- Needs
 - Prevention strategies
 - Role of PPIs, probiotics
 - Fidaxomicin in pediatric patients
 - Bezlotoxumab (Zinplava™)
 - New PCR-based laboratory platforms



References

1. McDonald LC, Gerding DN, Johnson S, Bakken JS, Carroll KC, Coffin SE, et al. Clinical practice guidelines for Clostridium difficile Infection in Adults and Children: 2017 update by the Infectious Diseases Society of America (IDSA) and Society for Healthcare Epidemiology of America (SHEA). *Clinical Infectious Diseases*. 2018;66(7):e1-e48.
2. Lawson PA, Citron DM, Tyrrell KL, Finegold SM. Reclassification of Clostridium difficile as Clostridioides difficile (Hall and O'Toole 1935) Prevot 1938. *Anaerobe*. 2016;40:95-9. Epub 2016/07/03. doi: 10.1016/j.anaerobe.2016.06.008. PubMed PMID: 27370902.
3. Bartlett JG, Chang TW, Gurwith M, Gorbach SL, Onderdonk AB. Antibiotic-associated pseudomembranous colitis due to toxin-producing clostridia. *N Engl J Med*. 1978;298(10):531-4. Epub 1978/03/09. doi: 10.1056/nejm197803092981003. PubMed PMID: 625309.
4. Mylonakis E, Ryan ET, Calderwood SB. Clostridium difficile--Associated diarrhea: A review. *Arch Intern Med*. 2001;161(4):525-33. Epub 2001/03/17. PubMed PMID: 11252111.
5. Young VB, Schmidt TM. Antibiotic-associated diarrhea accompanied by large-scale alterations in the composition of the fecal microbiota. *Journal of clinical microbiology*. 2004;42(3):1203-6.
6. Texts BL. 24.3: Bacterial Infections of the Gastrointestinal Tract 2017 [updated September 5, 2017July 6, 2018]. Available from: [https://bio.libretexts.org/TextMaps/Microbiology/Book%3A_Microbiology_\(OpenStax\)/24%3A_Digestive_System_Infections/24.3%3A_Bacterial_Infections_of_the_Gastrointestinal_Tract](https://bio.libretexts.org/TextMaps/Microbiology/Book%3A_Microbiology_(OpenStax)/24%3A_Digestive_System_Infections/24.3%3A_Bacterial_Infections_of_the_Gastrointestinal_Tract).
7. Matsuura H. Pseudomembranous colitis. *QJM: An International Journal of Medicine*. 2017;110(11):761-. doi: 10.1093/qjmed/hcx147.
8. Cohen SH, Gerding DN, Johnson S, Kelly CP, Loo VG, McDonald LC, et al. Clinical practice guidelines for Clostridium difficile infection in adults: 2010 update by the society for healthcare epidemiology of America (SHEA) and the infectious diseases society of America (IDSA). *Infection Control & Hospital Epidemiology*. 2010;31(5):431-55.
9. Magazine NG. Pufferfish 2018 [July 6, 2018]. Available from: <https://kids.nationalgeographic.com/animals/pufferfish/#pufferfish-closeup.jpg>.
10. Farooq PD, Urrunaga NH, Tang DM, von Rosenvinge EC. Pseudomembranous colitis. *Disease-a-month* : DM. 2015;61(5):181-206. Epub 2015/03/15. doi: 10.1016/j.disamonth.2015.01.006. PubMed PMID: 25769243; PubMed Central PMCID: PMC4402243.
11. Campus WG. What is PCR? 2016 [updated January 25, 2016July 6, 2018]. Available from: yourgenome.org/facts/what-is-pcr-polymerase-chain-reaction.
12. MyBioSource.com. What is ELISA? (Enzyme-linked Immunosorbent Assay). 2018.
13. Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *Journal of clinical epidemiology*. 2011;64(4):383-94. Epub 2011/01/05. doi: 10.1016/j.jclinepi.2010.04.026. PubMed PMID: 21195583.
14. Dellit TH, Owens RC, McGowan JE, Gerding DN, Weinstein RA, Burke JP, et al. Infectious Diseases Society of America and the Society for Healthcare Epidemiology of America guidelines for developing an institutional program to enhance antimicrobial stewardship. *Clinical infectious diseases*. 2007;44(2):159-77.
15. Post W. In fecal matter transplants, frozen is as good as fresh 2016 [updated January 12, 2016July 6, 2018]. Available from: https://www.washingtonpost.com/news/speaking-of-science/wp/2016/01/12/in-fecal-matter-transplants-frozen-is-as-good-as-fresh/?noredirect=on&utm_term=.c4d101d6f8d2.
16. Hoffmann D, Palumbo F, Ravel J, Roghmann M-C, Rowthorn V, von Rosenvinge E. Improving regulation of microbiota transplants. *Science*. 2017;358(6369):1390-1.
17. Staley C, Kelly CR, Brandt LJ, Khoruts A, Sadowsky MJ. Complete Microbiota Engraftment Is Not Essential for Recovery from Recurrent Clostridium difficile Infection following Fecal Microbiota Transplantation. *mBio*. 2016;7(6). Epub 2016/12/22. doi: 10.1128/mBio.01965-16. PubMed PMID: 27999162; PubMed Central PMCID: PMC45181777.
18. van Nood E, Vrieze A, Nieuwdorp M, Fuentes S, Zoetendal EG, de Vos WM, et al. Duodenal infusion of donor feces for recurrent Clostridium difficile. *N Engl J Med*. 2013;368(5):407-15. Epub 2013/02/27. doi: 10.1056/NEJMoa1205037. PubMed PMID: 23323867.

